



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

121. Proposed by W. J. GREENSTREET, M. A., Editor of The Mathematical Gazette, Stroud, Gloucestershire, England.

Two equal scale pans of mass m hang at rest over a smooth pulley. An inelastic particle, mass M , is dropped from a height h into one pan, and simultaneously another of equal mass and elasticity e is dropped from the same height into the other. Prove that every impact occurs when the pans are in their original positions, and find the total space described by either pan before motion ceases.

*** Solutions of these problems should be sent to B. F. Finkel not later than June 10.

AVERAGE AND PROBABILITY.

104. Proposed by LON C. WALKER, Assistant in Mathematics, Leland Stanford Jr. University, Palo Alto, Cal.

In a given sphere two radii are drawn at random, and a point taken in each at random. (1) Find the chance that the distance between the two points does not exceed the radius of the sphere. (2) Find the distance between them.

105. Proposed by LON C. WALKER, Assistant in Mathematics, Leland Stanford Jr. University, Palo Alto, Cal.

Find the average distance of the center of an ellipsoid, axes $2a$, $2b$, and $2c$, from its surface.

106. Proposed by LON C. WALKER, Assistant in Mathematics, Leland Stanford Jr. University, Palo Alto, Cal.

Required the average distance between two points in a hemisphere.

*** Solutions of these problems should be sent to B. F. Finkel not later than June 10.

MISCELLANEOUS.

105. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

If the refractive index of a medium at any point be $\mu = x$, prove that the path of the ray will be the curve $\frac{2x}{a} = \frac{c}{a} l^{y/a} + \frac{a}{c} l^{-(y/a)}$, a and c being constants.

106. Proposed by J. W. YOUNG, Oliver Graduate Student in Mathematics, Cornell University, Ithaca, N. Y.

Prove that $\frac{(2m)!}{(m!)^2}$ is an integer; and more generally that $\frac{(nm)!}{(m!)^n}$ is an integer, m , n being any positive integers.

107. Proposed by WILLIAM HOOVER, A.M., Ph.D., Professor of Mathematics and Astronomy, Ohio University, Athens, Ohio.

The index of refraction of a medium varying inversely as the square root of the distance, prove that the path of a ray of light in the medium is a cycloid.

*** Solutions of these problems should be sent to J. M. Colaw not later than June 10.